Deepwater Program: Exploration and Research of Northern Gulf of Mexico Deepwater Natural and Artificial Hard Bottom Habitats with Emphasis on Coral Communities: Reefs, Rigs and Wrecks

Project Director - James Brooks

TDI-Brooks International, 1902 Pinon, College Station, TX 77845 Phone: (979) 693-3446 FAX: (979) 693-6389 E-mail: <u>Drjmbrooks@aol.com</u>

CO-PI Chuck Fisher

208 Mueller Laboratory ,The Pennsylvania State University, University Park, PA 16802 Phone: (814) 865-3365 FAX: (814) 865-9131 E-mail: <u>cfisher@psu.edu</u>

CO-PI Harry Roberts

CSI-LSU, 304 Howe-Russell Geosciences Complex, Baton Rouge, LA 70803 Phone: 225.578-2964 FAX: 225-578-2520 E-mail: hrober3@lsu.edu

CO-PI Dan Warren, Robert Church C&C Technologies, Inc, 730 E. Kaliste Saloom Rd., Lafayette, LA 70508 Phone: 337-261-0660 E-mail: djw@cctechnol.com, rc@cctechnol.com

CO-PI Ian MacDonald Florida State University, Rm 511 OSB, Tallahassee, FL 32306-4320 Phone: (850) 644-5498 FAX: (850) 644-2581 E-mail: imacdonald@fsu.edu

CO-PI Erik Cordes

315 Biology-Life Sciences, 1900 N. 12th Street, Temple University, Philadelphia, PA 19122
Phone: 215-204-8854 FAX: 215-204-6646 E-mail: erik.cordes@temple.edu

CO-PI Chris German

Woods Hole Oceanographic Institution, Clark South 276, MS#24, Woods Hole, MA 02543 Phone: 508 289 2853 FAX: E-mail: cgerman@whoi.edu

CO-PI Timothy M. Shank

240 Redfield Laboratory, Woods Hole Oceanographic Institution, Woods Hole, MA 02543 Phone: 508 289 3392 FAX: E-mail: tshank@whoi.edu

CO-PI Liz Goehring NEON, Inc.,1685 38th St., Ste. 100, Boulder, CO 80301 Phone: (720) 746-4844 FAX: (720) 746-4870 E-mail: lgoehring@neoninc.org

CO-PI Bernie Bernard TDI-Brooks International, 1902 Pinon, College Station, TX 77845 Phone: 979.690-6287 FAX: (979) 693-6389 E-mail: <u>berniebernard@tdi-bi.com</u>

CO-PI Gary Wolff TDI-Brooks International, 1902 Pinon, College Station, TX 77845 Phone: 979.846-7679 FAX: (979) 693-6389 E-mail: <u>garywolff@geodatapub.com</u>

> Award Number: M08PC20038 http://www.tdi-bi.com/Lophelia/lophelia-main.htm

LONG-TERM GOALS

A primary goal of this study is to obtain a robust predictive capability for the occurrence of rich cnidarian (primarily scleractinian coral) hard ground communities in the deep Gulf of Mexico (GoM). To achieve this long-term goal, this study will accomplish three interrelated and interdependent objectives.

OBJECTIVES

The objectives of this study are:

- 1) Discover and describe new locations at greater than 300m depth in the GoM with extensive coral community development, particularly including *Lophelia pertusa*.
- 2) Gain a more comprehensive understanding of the fundamental processes that control the occurrence and distribution of *Lophelia* and other extensive coral communities at depths greater than 300 m in the GoM through both laboratory experiments and field data collection.
- 3) Document and understand the relations between coral communities on artificial and natural substrates with respect to community composition and function, phylogeographic and population genetics, and growth rates of the key cnidarian foundation fauna.

Our specific biological objectives:

- 1) To discover and characterize new sites:
 - a. Characterize key sites at the largest scale with high resolution bathymetry, side scan sonar imaging, 3D seismic data, and current models
 - b. Characterize the coral density at all study sites at the 10 to 100 m scale with randomized photo transects and general site descriptions.
 - c. Characterize the community composition at the 1 to 10 m scale at significant coral sites (man-made and natural) with analysis of close up imagery, replicate 5x5m photomosaics, and quantitative community collections.
- 2) Analyze connectivity among all sites in the GoM (man-made and natural), and to other sampled populations, with comparative community, phylogeographic and population genetic analyses.
- 3) Compare the structure, species richness and diversity of communities tightly associated with Lophelia on man-made structures and from natural sites in the GoM
- 4) Experimentally determine the tolerance and growth response of Lophelia to temperature, pH/alkalinity, dissolved oxygen and electrical current.
- 5) Characterize and constrain growth rates of key species of colonial cnidarians (pioneer colonies) using analyses of images of the largest colonies on man-made structures of known age.
- 6) Characterize key variables at sites with the most significant communities of coral colony development over one year by monitoring temperature, currents, larval seasonal distribution and sediment quality and quantity at 2-4 sites between 2009 and 2010.

By integrating this information, we will develop a predictive model to examine the potential occurrence of significant assemblages of *L. pertusa* or other cnidarians at unexplored sites and artificial reefs in the deep GoM.

Historic Shipwreck Component

The proposed study will be multidisciplinary in scope. It will focus on the archaeological and biological aspects of up to six shipwrecks in the north-central portion of the GoM. Water depths at the designated investigation sites range from 554m to 2,286m. Each shipwreck will be investigated to

determine identity, site boundaries, National Register eligibility, preservation state and stability, associated biological communities, and an artificial reef effect on meiofauna and/or mobile fish communities.

Our specific archaeological objectives:

1) To record each vessel through detailed imagery to establish its type, date of construction, and positive identification if possible.

2) To establish nationality, ownership (past and present), use history, cause of loss, mission and cargo at time of loss through fieldwork and historical research.

3) To determine the extent and condition of the artifact assemblage on each vessel and the presence of diagnostic artifacts.

4) To determine potential eligibility to the National Register of Historic Places through archival research and the analysis of imagery and to prepare a National Register nomination form for potentially eligible vessels.

5) To assess impacts of biofouling communities to these shipwrecks to determine the stability of these sites and rate of deterioration.

Our specific bacteriological objectives:

1) Determine the rates of deterioration of test coupons placed on platforms and sited on specific shipwrecks that have already been visited and were installed on previous expeditions to those sites of interest. Determination may take the form of passive observations of the coupon condition within the platform of interest. If significant deterioration rates have already occurred then the platform may be recovered for on-ship and in-laboratory investigations.

2) Fresh test platforms may be deployed to either replace recovered platforms or to determination the deterioration rate at a fresh site of archeological significance.

3) Examine recovered natural samples from deteriorating sites and localized waters that may contain active bacterial communities traditionally associated with corrosive or bio-concretious growths.

4) On selected recovered test coupons conduct such experiments as would allow a more precise projection of the rates of deterioration of test material through the rates of pitting (in the event of mild steel) to losses in physical composition (in the event of cloths and wood.

5) Recovered bacteriological communities from the sites of interest will be subjected to laboratory analysis to determine major bacterial communities, degrees of activity (using the ATP methodology), and identification of community structures by fatty acid methyl ester analysis (MIDI/BART technologies).

APPROACH AND WORK PLAN

In order to meet the objectives outlined above, the following scientific and technical plan is being implemented.

Key individuals participating in this work and their roles are: Dr. James Brooks is the Project Manager and takes the lead in administration of this project. Dr. Charles Fisher (Pennsylvania State University)

coordinates the biological studies, Dr. Harry Roberts (Louisiana State University) coordinates the geological/geophysical and oceanographic studies, Mr. Dan Warren and Mr. Rob Church from C&C Technology coordinate the wrecks studies along with, Dr. Chris German (WHOI) who coordinates the time series sediment trap studies, and Ms. Liz Goehring (NEON and NSF Ridge 2000 office) coordinates the education and outreach activities.

Dr. Charles Fisher oversees the biological aspects of the study and the interface with the USGS teams and the geological and oceanographic measurements and studies. He works closely with Dr. Roberts, Dr. Cordes, Dr. German, and other PIs to plan and conduct the submersible/ROV portions of the field work. His research group will take responsibility for quantitative physical and photographic collections, community composition and structure analyses, and trophic studies of the endemic and other closely associated seep and coral fauna. Dr. Harry Roberts takes the lead in site selection for exploration of potential new natural deep coral communities and also coordinates the geological and oceanographic deployments and studies. His colleague, Dr. Susan Welsh develops and runs a very high resolution numerical circulation model of the Gulf of Mexico that will be used to help identify the likely geographic areas and depth ranges for recruitment of L. pertusa larvae on the northern slope and compute dispersal kernel trajectories and dispersion of coral larvae by oceanographic current. Dr. Erik Cordes takes responsibility for the live Lophelia pertusa experiments, gorgonian genetics, and coordinate the phylogeography and taxonomy team. He also works with Fisher's team on studies of coral communities and is responsible for coordination with other outside taxonomic experts. Dr. Ian MacDonald directs the use of digital imagery in the initial site survey and recon cruise, and deployment and analyses of the time lapse cameras and imagery. Dr. Chris German takes the lead on deployments of the time series sediment traps and oversee the integrated analyses of the samples. Dr. William Schroeder assists in the analysis of coral communities on rigs and wrecks and also participates in the over-all study synthesis phase. Dr. Tim Shank conducts studies on the population genetics of several coral associated taxa. Dr. Iliana Baums oversees development of molecular markers for studies of population connectivity in antipatharians and also serve as consultant for the construction of live Lophelia maintenance aquaria. Dr. Stephen Cairns is responsible for the identification of the hard coral samples collected. Peter Etnoyer oversees video and the classical taxonomy of octocorals and Dr. Dennis Opresko the classical taxonomy and phylogeography of antipatharians.

In addition to this core team we have assembled an international team of collaborators that significantly expands our taxonomic expertise. Limpets and snails are sent to Anders Waren (Swedish Museum of Natural History) for morphological characterization. Dr. Stéphane Hourdez (Statione Biologique de Roscoff, France) takes the lead on polychaete phylogenetic characterizations and descriptions of new species (using both molecular and classical approaches). Dr. Daphne Fautin (University of Kansas) assists in the taxonomic identification of sea anemones Dr. Sabine Stohr (Swedish Museum of Natural History) examines all new ophuiroids collected. Maria Pia Miglietta (Penn State University) leads the morphological and genetic identification of the hydroids.

C & C Technologies, Inc. (C & C) of Lafayette, LA has partnered with Droycon Bioconcepts Inc. (DBI) of Regina Canada to fulfill the archaeological component of the study. Marine archaeologists Daniel Warren and Robert Church are the C & C project managers and co-principle investigators. Dr. Sheli Smith is the director of operations for the PAST Foundation, a subcontractor to C & C. She has substantial knowledge in eighteenth and nineteenth century marine architecture and is assessing three to four of the wreck sites in this study from those periods.

Dr. Roy Cullimore, the CEO of DBI, and Lori Johnston primary role on this project is to evaluate the deterioration rate, preservation, and stability of the wreck sites. Re-examining test platforms placed during the earlier studies will be instrumental to this analysis. They coordinate with Dr. Christina A. Kellogg with USGS for other microbiology aspects during the project.

Dr. Bernie Bernard, TDI-Brooks Director and Chief Technology Officer, coordinates the navigation and logistics portion of the project. Dr. Gary Wolff is the project's Data Manager as he has been for other BOEM projects including the Chemosynthesis-series. Mrs. Gail Mills provides financial and project administrative support.

WORK COMPLETED

Four cruises have been completed since Lophelia II began. **Table** 1 summarizes the dates, ships used and objectives.

Cruise	Dates	Ship	Comments
1	2 September - 2 October	Nancy Foster - NOAA	Ewing Banks Wreck, Gulfoil,
	2008		Gulfpenn sites studied
2	16 June - 1 July 2009	RV Brooks McCall – TDI-	multibeam collected at 13
		Brooks	sites and ROV at 8 sites
3	19 August - 12 September	Ron Brown - NOAA	multibeam swath bathymetry,
	2009		digital video and photographs
4	13 October - 4 November,	Ron Brown - NOAA	17 ROV dives including the
	2010		Macondo well area

RESULTS

Cruise1 Leg1 – Using NOAA Ship *Nancy Foster / SeaVision* ROV, the Ewing Banks Wreck site was confirmed to be a historic shipwreck site, which likely dates to the nineteenth century. More *Lophelia* was discovered growing on that site than any other nineteenth century wooden wreck known in the Gulf of Mexico. Many of the wreck's attributes, such as the lack of cargo, rigging, or machinery, make it an intriguing mystery. The identity of *Gulfoil* was confirmed on a brief dive at the site and substantial coral colonies were documented. The preliminary examination suggests the *Lophelia* coverage at *Gulfoil* may be more substantial than that documented at *Gulfpenn* in 2004. The dive on *Gulfpenn* allowed the identification of the stern section of the tanker, the microbial experiment placed on the site in 2004 was reexamined, and a temperature logger was placed on *Gulfpenn*'s bow.

Cruise1 Leg2 - During the 12 working days at sea, multibeam data was collected at 13 sites and 10 lowerings of the ROV were completed over 8 different sites. There were two ROV operations days lost to weather conditions, one dive never reached the bottom, and three other dives were ended with under two hours of bottom time due to technical problems with the ROV. During almost every dive the manipulator failed, and this severely limited the capacity to collect physical samples for future genetics work. On most of the latter dives, only one attempt was made to collect samples in areas of high coral diversity and density due to the anticipation of manipulator failure and the extremely long amount of time required to sample (1-5 samples obtained in 1-3 hours).

Cruise 2 - The cruise team was at sea for 15 days on board the *RV Brooks McCall* and transited from Freeport, TX to the Garden Banks Lease Area south of the Flower Garden Banks and then to the Green

Canyon lease area. The primary data collected includes multibeam swath bathymetry, digital video and still photographic imagery, and CTD with DO and pH sensors. Other data streams from the AUV, such as vehicle attitude, acoustic data, and sonar imagery are recorded by networked computers in the control van. Navigational data for both the ship and AUV systems will also be recorded. While in transit to and from the site, and during times when the AUV is not deployed, photographic data will be collected with the camera sled.

Cruise 3 - *Jason II* was used to: explore 10 new sites for the occurrence of deep water coral reefs; make collections of *Lophelia* and other corals for genetic and physiological studies, make collections of communities associated with *Lophelia* and other corals for ecological studies; collect quantitative digital imagery for characterization of sites and coral communities; collect spatially explicit physical near bottom oceanographic data; deploy cameras and microbial arrays; reposition larval traps and current meters; collect push cores; and conduct a series of linked archeological/biological investigations on deep water shipwrecks. In addition to launching and recovering *Jason II*, elevators were deployed and recovered twice, four moorings (2 larval traps and 2 current meters) were deployed, and CTD casts were conducted. 19 August – 12 September 2009

Cruise 4 - The cruise was completed on NOAA Ship *Ronald H. Brown* from 13 October – 4 November 2010. The cruise mobilized in Pensacola, Florida. One mid-cruise personnel transfer took place on 24 October. The cruise demobilized in Pensacola, Florida on 5 November 2010. This was a 22-day cruise with 17 ROV dives including the Macondo well area.

IMPACT AND APPLICATIONS

National Security

This program will provide critical information on the exploration, characterization and function of hardground communities to BOEM. As manager of the nation's seafloor mineral resources, BOEM will use this information to aid in the development of critical energy resources, which may affect domestic energy production.

Economic Development

Increased energy and mineral production will have a positive economic impact at numerous levels in industry.

Quality of Life

Information on the location and functioning of hardground communities gathered by this program will have a positive impact on other ocean users, the natural environment, and the human environment. It will aid in minimizing the environmental impact on sensitive habitat and mitigate any potential damage to these communities.

Science Education and Communication

Education outreach efforts outside of the cruise website build on and leverage from the success of both the NOAA OE educational materials and the existing FLEXE (From Local to Extreme Environments) educational project. Full instructional units have been developed on deep-sea coral ecology integrated with the NOAA/OE lessons, multimedia modules and Ocean Explorer website featured as the GLOBE FLEXE campaign for the 2010-2012 academic years.

TRANSITIONS

Data is provided to the ChEss database, which is a component of the Census of Marine Life (CoML) Ocean Biogeographic Information System (OBIS) data base. This work will contribute significantly to the goals of the Atlantic Equatorial Belt studies of the ChEss program, particularly the components that will allow interpretation of our findings in the context of hardground habitats around the world.

RELATED PROJECTS

- Chemosynthetic Ecosystems Study (MMS Report 95-0021). <u>http://www.gomr.mms.gov/homepg/regulate/environ/studies/1995/95-0021%20Vol%20I.pdf</u>.
- Stability and Change in Gulf of Mexico Chemosynthetic Communities (MMS Report 2002-036). <u>http://www.gomr.mms.gov/homepg/regulate/environ/studies/2002-036.pdf</u>.
- The Deepwater Program: Northern Gulf of Mexico Continental Slope Habitat and Benthic Ecology (MMS contract 1435-01-99-CT-30991).
- Investigations of Chemosynthetic Communities on the Lower Continental Slope of the Gulf of Mexico. (Award Number: 1435-01-05-39187) <u>http://tdi-bi.com/chemo3/chemo-main.htm</u>



Figure 1. View of the Lophelia colony growing on the EW Wreck's stem post.



Figure 2. Station locations of the project's four cruises.

REFERENCES RELATED TO THIS PROJECT

- Aloisi, G., I. Bouloubassi, S.K. Heijs, R.D. Pancost, C. Pierre, J.S.S. Damste, J.C. Gottschal, L.J. Forney and J.M. Rouchy, 2002, CH₄-consuming microorganisms and the formation of carbonate crusts at cold seeps: Earth and Planetary Science Letters, v. 203, p. 195-203.
- Andrews A.H., E.E. Cordes, M.M. Mahoney, K. Munk, K.H. Coale, G.M. Cailliet, J. Heifetz. 2002. Age, growth and radiometric age validation of a deep-sea, habitat-forming gorgonian (*Primnoa resedaeformis*) from the Gulf of Alaska. <u>Hydrobiologia</u> 471: 101-110.
- Barnette, M. C. 2001. A review of the fishing gear utilized within the Southeast Region and their potential impacts on essential fish habitat. NOAA Tech. Memo. NMFSSEFSC- 449. 62 pp.
- Barnette, M. C. 2006. Observations of the Deep-Water Coral Oculina varicosa in the Gulf of Mexico
- Baums IB, Hughes CR, Hellberg MH (2005a) Mendelian microsatellite loci for the Caribbean coral *Acropora palmata. Marine Ecology Progress Series* **288**, 115-127.
- Baums IB, Miller MW, Hellberg ME (2005b) Regionally isolated populations of an imperiled Caribbean coral, *Acropora palmata*. *Molecular Ecology* **14**, 1377-1390.
- Baums IB, Paris CB, Cherubin LM (2006) A bio-oceanographic filter to larval dispersal in a reefbuilding coral. *Limnology and Oceanography* **51**, 1969-1981.
- Bayer, FM (1956) Octocorallia. In: Moore, RC (ed) Treatise on invertebrate paleonotology. Part F, Coelenterata. University of Kansas Press, Lawrence, Kansas, p 166-231
- Berman, Bruce D. 1971. Encyclopedia of American Shipwrecks. Boston: Mariners Press.
- Bergquist, D. C., I. A. Urcuyo, and C. R. Fisher. 2002. Establishment and persistence of seep vestimentiferan aggregations from the upper Louisiana slope of the Gulf of Mexico. <u>Mar. Ecol.</u> <u>Prog. Ser</u>. 241: 89-98.
- Bett BJ (1997) Atlantic Margin Environmental Survey: Seabed Survey of the Shelf Edge and Slope west of Shetland. Published by Challenger Division for Seafloor Processes: Southampton Oceanography Centre, NERC/ University of Southampton., pp. 165.
- Bett B.J. (2001). UK Atlantic margin environmental survey: introduction and overview of bathyal benthic ecology. *Cont. Shelf Res.* 21: 917-956.
- Browning, Robert M., Jr. 1996. U.S. Merchant Vessel War Casualties of World War II. Naval Institute Press. Annapolis.
- Burch, H. A. 1942a. Summary of Statements by Survivors of the SS *Gulf Penn*, U.S. Tanker. Navy department, Office of the Chief of Naval Operations. May 22. NOAA Technical Memorandum NMFS-SEFSC-535. 12 pp.
- Cairns, SD (1977) Guide to the commoner shallow water gorgonians (sea whips and sea fans) of Florida, the Gulf of Mexico, and the Caribbean. Sea Grant Field Guide Ser. No. 6

- Cairns S.D. (1978). A checklist of the ahermatypic scleractinia of the Gulf of Mexico, with the description of a new species. *Gulf Res. Rep.* 6: 9-15.
- Cairns S.D. (2001). A brief history of taxonomic research on azooxanthellate Scleractinia (Cnidaria: Anthozoa). *Bull. Biol. Soc. Wash.* 10: 191-203.
- Cairns S.D., D.M. Opresko, T.S. Hopkins & W.W. Schroeder. (1993). New records of deep-water Cnidaria (Scleractinia & Antipatharia) from the Gulf of Mexico. *Northeast Gulf Sci.* 13:1-11.
- Childress, J.J. and T.J. Mickel. 1980. A motion compensated shipboard precision balance system. Deep-Sea Res. 27: 965-970.
- Church, R.A. 2003. Archaeological assessment for a proposed 6-inch gas and 6-inch oil pipeline routes from Block 962 to Block 786, Viosca Knoll Area. Prepared by C & C Technologies, Inc. for Geoscience Earth & Marine Services, Inc. on behalf of Mariner Energy. Report submitted to the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Church, R., D. Warren, R. Cullimore, L. Johnston, W. Schroeder, W. Patterson, T. Shirley, M. Kilgour, N. Morris, and J. Moore. 2007. Archaeological and biological analysis of World War II shipwrecks in the Gulf of Mexico: Artificial reef effect in deep water. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2007-015.
- Church, Robert. 2004. "Shipwreck and Sonar Contact ROV Investigation, Blocks 245 and 329, Green Canyon Area." Prepared by C & C Technologies, Inc. for GulfTerra Energy Partners, L.P. Report submitted to the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Church, R.A. and D.J. Warren. 2008. Viosca Knoll Wreck: Discovery and investigation of an early nineteenth-century sailing ship in 2,000 feet of water. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2008-018.
- Church, Robert and Lynn Samuel. 2004. Archaeological, Engineering and Hazard Study for a Proposed 16-inch Oil Export Pipeline from Block 680, Green Canyon Area to Block 332, Ship Shoal Area. Prepared by C & C Technologies, Inc. for GulfTerra Energy Partners, L.P. Report submitted to the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Church, R., D. Warren, R. Cullimore, L. Johnston, W. Schroeder, W. Patterson, T. Shirley. M. Kilgour, N. Morris, and J. Moore. 2007. <u>Archaeological and Biological Analysis of World War II</u> <u>Shipwrecks in the Gulf of Mexico: Artificial Reef Effect in Deep Water</u>. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2007-015. 387 pp.
- Clarke, K. R. and R. M. Warwick (2001). Change in marine Communities: an approach to statistical analysis and interpretation. Plymouth, England, PRIMER-E.

Colin P (1975) The neon gobies T. F. H. Publ., Neptune City, NJ.

- Cordes, E.E., Bergquist, D.C., Predmore, B.L., Dienes, P., Jones, C., Telesnicki, G., Fisher, C.R., 2006. Alternate unstable states: convergent paths of succession in hydrocarbon-seep tubewormassociated communities. Journal of Experimental Marine Biology and Ecology 339, 159-176.
- Cordes EE, McGinley M, Podowski EL, Becker, EL, Lessard-Pilon S, Viada S, Fisher CR. (2008) Coral communities of the deep Gulf of Mexico. Deep-Sea Research I. 10.1016/j.dsr.2008.03.005
- Continental Shelf Associates, Inc. 2002. Deepwater Program: Bluewater Fishing and Deepwater OCS Activity, Interactions between the Fishing and Petroleum Industries in Deepwaters of the Gulf of Mexico. A Final Report for the U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, Metairie, LA. OCS Study MMS-2002-78.
- Continental Shelf Associates International, Inc. 2007. Characterization of northern Gulf of Mexico deepwater hardbottom communities with emphasis on *Lophelia* coral. U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, Metairie, LA. OCS Study MMS-2007-44.
- Cowen RK, Paris CB, Srinivasan A (2006) Scaling of connectivity in marine populations. *Science* **311**, 522-527.
- Davies, P. S. (1989) Short-term growth measurements of corals using an accurate buoyant weighing technique Mar. Biol. Vol. 101: 389-395
- Dodds LA, Roberts JM, Taylor AC, Marubini F. 2007. Metabolic tolerance of the cold-water coral Lopehlia pertusa (Scleractinia) to temperature and dissolved oxygen change. Journal of Experimental Marine Biology and Ecology. 349: 205-214.
- Eslinger, Kimberly. 2006. "Cultural Resources Assessment Report, "Medusa Prospect", Blocks 493-496, 537-540, 581-583, 626-627, and Vicinity, Mississippi Canyon Area." Prepared by C & C Technologies, Inc. for Murphy Exploration & Production Company. Report submitted to the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Etnoyer, P. and J. Warrenchuk (2007). A catshark nursery in a deep gorgonian field in the Mississippi Canyon, Gulf of Mexico. Bulletin of Marine Science 81: 553-559.
- Formolo, M.J., Lyons, T.W., Zhang, C., Kelley, C., Sassen, R., Horita, J., Cole, D.R., 2004. Quantifying carbon sources in the formation of authigenic carbonates at gas hydrate sites in the Gulf of Mexico. Chemical Geology 205, 253-264.
- Fossa, J. H., Mortensen, P. B., Furevik, D. M. 2002. The deep-water coral *Lophelia pertusa* in Norwegian waters: distribution and fishery impacts. Hydrobiologia 471:1-12.
- Frederiksen R, Jensen A, and Westerberg H (1992) The distribution of the scleractinian coral *Lophelia pertusa* around the Faroe Islands and the relation to internal mixing. Sarsia 77:157-171.

- Freiwald, A., R. Henrich, J. Patzold. 1997. Anatomy of a deep-water coral reef mound from Stjernsund, west Finnmark, northern Norway. In James, N. P., Clarke, J. A. D., (eds.), Cool-Water Carbonates. Society of Sedimentary Geologists (SEPM), Special Publication. 56:141-162.
- Freiwald, A., Henrich, R., Patzold, J. 1997. Anatomy of a deep-water coral reef mound from Stjernsund, west Finnmark, northern Norway. *In James*, N. P., Clarke, J. A. D., eds., Cool-Water Carbonates. Society of Sedimentary Geologists (SEPM), Special Publication. 56:140-161.
- Freiwald A (1998) Geobiology of *Lophelia pertusa* (Scleractinia) reefs in the North Atlantic. Habilitation, University of Bremen.
- Freiwald A., J.B. Wilson & R. Henrich. (1999). Grounding Pleistocene icebergs shape recent deepwater coral reefs. *Sed. Geol.* 125: 1-8.
- Freiwald, A., V. Huhnerbach, B. Lindberg, J.B. Wilson, and J. Campbell. 2002. The Sula Ridge Complex, Norwegian Shelf. Facies 47:179-200.
- Gass SE, Roberts JM (2006) The occurrence of the cold-water coral Lophelia pertusa (Scleractinia) on oil and gas platforms in the North Sea: colony growth, recruitment and environmental controls on distribution. Marine Pollution Bulletin 52: 549-559
- Hall-Spencer, J., Allain, V., Fossa, J. H. 2002. Trawling damage to Northeast Atlantic ancient coral reefs. Proc. R. Soc. Lond. B. 269:507-511.
- Harper, D. E., P. B. Eyo and G. P. Scott. 2000. Updated golden crab fishery trends and production model analysis based on trip report logbook and trip interview data collection programs. Contribution Number PRD-99/00-12. NMFS Southeastern Fisheries Science Center. Miami, Florida.
- Hayes, J.A. 1990. Distribution, movement and impact of the corallivorous gastropod *Coralliophila abbreviata* (Lamark) on a Panamanian patch reef. J. Exp. Mar. Biol. Ecol. 142:25-42.
- Heifetz J. (2002) Coral in Alaska: distribution, abundance, and species associations. *Hydrobiologia* 471: 19-28.
- Hellberg ME (2006) No variation and low synonymous substitution rates in coral mtDNA despite high nuclear variation. *Bmc Evolutionary Biology* **6**. *Evolution* **53**, 806-817.
- Henry, L-A., Roberts, J.M., 2007. Biodiversity and ecological composition of macrobenthos on coldwater coral mounds and adjacent off-mound habitat in the bathyal Porcupine Seabight, NE Atlantic. Deep-Sea Res. I. 54 (4), 654–672.
- Horrell, Christopher E. 2006. "500 Fathoms of Water: The Archaeological Remains of a Mid-Late Nineteenth Century Sailing Vessel in the Gulf of Mexico." *Proceedings of the Society for Historical Archaeology Conference*, Sacramento, CA (January).
- Hovland, M., Thomsen, E. 1997. Cold-water corals- are they hydrocarbon seep related? Mar. Geol. 137: 159-164.

- Hovland, M., Mortensen, R. B., Brattegard, T., Strass, P., Rokoengen, K. 1998. Ahermatypic coral banks off mid-Norway: Evidence of a link with seepage of light hydrocarbons. Palaios 13:189-200.
- Jensen A., and R. Frederiksen. 1992. The fauna associated with the bank-forming deepwater coral *Lophelia pertusa* (Scleractinia) on the Faroe shelf. Sarsia 77:53-69.
- Kleypas, J.A., R.A. Feely, V.J. Fabry, C. Langdon, C.L. Sabine, and L.L. Robbins, 2006. Impacts of Ocean Acidification on Coral Reefs and Other Marine Calcifiers: A Guide for Future Research, report of a workshop held 18–20 April 2005, St. Petersburg, FL, sponsored by NSF, NOAA, and the U.S. Geological Survey, 88 pp.
- Krieger KJ, Wing BL (2002) Megafauna associations with deepwater corals (*Primnoa* spp.) in the Gulf of Alaska. Hydrobiologia 471: 82-90
- Koslow J.A., K. Gowlett-Holmes, J.K. Lowry, T. O'Hara, G.C.B. Poore, and A. Williams. (2001). Seamount benthic macrofauna off southern Tasmania: community structure and impacts of trawling. Mar. Ecol. Prog. Ser. 213:111-125.
- Lockhart, F. D., W. J. Lindberg, N. J. Blake, R. B. Erdman, H. M. Perry, and R. S. Waller. 1990. Distributional differences and population similarities for two deep-sea crabs (Family Geryonidae) in the northeastern Gulf of Mexico. Canadian Journal of Fisheries and Aquatic Sciences 47: 2112-2122.
- Loya Y. and B. Rinkevich. (1980). Effects of oil pollution on coral reef communities. *Mar. Ecol. Prog. Ser.* 3: 167-180.
- Masson D.G, B.J. Bett, D.S.M. Billett, C.L. Jacobs, A.J. Wheeler and R.B. Wynn. (2003). The origin of deep-water, coral-topped mounds in the northern Rockall Trough, Northeast Atlantic. *Mar. Geol.* 194: 159-180.
- McEachran J.D. and J.D. Fechhelm. (1998). *Fishes of the Gulf of Mexico*. University of Texas Press, Austin, TX. 1112 pp.
- Mienis, F., H. C. de Stigter, M. White, G. Dulneveldc, H. de Haas and T. C. E. van Weering (2007). "Hydrodynamic controls on cold-water coral growth and carbonate-mound development at the SW and SE rockall trough margin, NE Atlantic ocean." Deep-Sea Research Part I-Oceanographic Research Papers 54(9): 1655-1674.
- Mikkelsen N, Erlenkauser H, Killingley JS, and Berger WH (1982) Norwegian corals: radiocarbon and stable isotopes in *Lophelia pertusa*. Boreas 5:163-171.
- Moore, D. R., Bullis, H. R. 1960. A deep-water coral reef in the Gulf of Mexico. Bulletin of Marine Science. Vol. 10:125-128
- Mortensen, P. B., Hovland, M., Brattegard, T., Farestveit, R. 1995. Deep-water bioherms of the scleractinian coral *Lophelia pertusa* L. at 64° on the Norwegian shelf: structure and associated megafauna. Sarsia. 80:145-158.

- Mortensen, P. B., Rapp, H. T. 1998. Oxygen and carbon isotope ratios related to growth line patterns in skeletons of *Lophelia pertusa* (L) (Anthozoa, scleractinia): implications for determinations of linear extension rates. Sarsia. 83:433-446.
- Neigel JE (1997) A comparison of alternative strategies for estimating gene flow from genetic markers. *Annual Review of Ecology and Systematics* **28**, 105-128.
- Neumann AC, Kofoed JW, and Keller GH (1977) Lithoherms in the Straits of Florida. Geology 5:4-10.
- Penn K, Wu D, Eisen JA, Ward N (2005) Characterization of Bacterial Communities Associated with Deep-Sea Corals on Gulf of Alaska Seamounts. App Env Microbiol 72(2): 1680-1683
- Perry, H., R. Waller, C. Trigg, J. McBee, R. Erdman, and N. Blake. 1995. A note on bycatch associated with deepwater trapping of *Chaceon* in the northcentral Gulf of Mexico. Gulf Research Reports 9(2): 139-142.
- Peterson C.H., M.C. Kennicutt, R.H. Green, P. Montagna, D.E. Harper, E.N. Powell and P.F. Roscigno. (1996). Ecological consequences of environmental perturbations associated with offshore hydrocarbon production: a perspective on long-term exposure in the Gulf of Mexico. *Can. J. Fish. Aquat. Sci.* 53: 2637-2654.
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* **155**, 945-959.
- Reed J.K. (2002). Comparison of deep-water reefs and lithoherms off southeastern USA. *Hydrobiologia* 471: 57-69.
- Reed, J. K., D. C. Weaver, and S. A. Pomponi. 2006. Habitat and fauna of deep-water *Lophelia pertusa* coral reefs off the southeastern U.S.: Blake Plateau, Straights of Florida, and Gulf of Mexico. Bulletin of Marine Science 78(2): 343-375.
- Reed, J. K., S. A. Pomponi, D. C. Weaver, C. K. Paull, and A. E. Wright. 2005. Deep-water sinkholes and bioherms of South Florida and the Pourtales Terrace- Habitat and Fauna. Bulletin of Marine Science 77(2):267-296.
- Risk M.J., J.M. Heikoop, M.G. Snow, R. Beukens. (2002). Lifespans and growth patterns of two deepsea corals: *Primnoa resedaeformis* and *Desmophyllum cristagalli*. *Hydrobiologia* 471: 125-131.
- Ritger, S., B. Carson, and E. Suess, 1987, Methane-drived authigenic carbonates formed by subduction-induced pore-water expulsion along the Oregon/Washington margin: Geological Society of America Bulletin, v. 98, p. 147-156.
- Roberts, H.H. and P. Aharon, 1994, Hydrocarbon-derived carbonate buildups of the northern Gulf of Mexico continental slope: A review of submersible investigations: Geo-Marine Letters, v. 14, p. 135-148.
- Roberts, H.H, B.A. Hardage, W.W. Shedd, and J. Hunt, Jr., 2006, Seafloor reflectivity An important seismic property for interpreting fluid/gas expulsion geology and the presence of gas hydrate: The Leading Edge, Society of Exploration Geophysicists, v. 25, p. 620-628.

- Roberts HH, Fisher CR, Bernard B, Brooks JM, Bright M, Carney RS, Cordes EE, Goehring L, Hourdez S, Hunt Jr. JL, Joye SM, MacDonald IR, Morrison C, Niemann H, Potter J, Samarkin V, Shedd W, Telesnicki G, Becker EL, Bernier M, Bowles M, Kupehik M, Lessard-Pilon S, Petersen C, Petersen J. (2007) *ALVIN* explores the deep northern Gulf of Mexico slope. EOS 88: 341-342.
- Roberts JM, and Anderson RM (2002) A new laboratory method for monitoring deep-water coral polyp behaviour. Hydrobiologia 471:143-148.
- Roberts J.M., D. Long, J.B. Wilson, P.B. Mortensen, J.D. Gage. (2003). The cold-water coral *Lophelia pertusa* (Scleractinia) and enigmatic seabed mounds along the north-east Atlantic margin: are they related? *Mar. Poll. Bull.* 46: 7-20.
- Roberts, J. M. (2005). "Reef-aggregating behaviour by symbiotic eunicid polychaetes from cold-water corals: do worms assemble reefs?" Journal of the Marine Biological Association of the United Kingdom 85(4): 813-819.
- Roberts, J.M., Wheeler, A.J., Freiwald, A., 2006. Reefs of the deep: The biology and geology of coldwater coral ecosystems. Science 312, 543-547.
- Rogers, A. D. 1999. The biology of *Lophelia pertusa* (Linnaeus 1758) and other deep-water reef forming corals and impacts from human activities. Int. Rev. Hydrobiol. 844:315-406.
- Rosenberg NA, Burke T, Elo K, *et al.* (2001) Empirical evaluation of genetic clustering methods using multilocus genotypes from 20 chicken breeds. *Genetics* **159**, 699-713.
- Ross, S. W. and A. M. Quattrini. 2007. The fish fauna associated with deep coral banks off the southeastern United States. Deep-Sea Research I 54: 975-1007.
- Sabater MG and Yap HT (2002) Growth and survival of coral transplants with and without electrochemical deposition of CaCO3. Journal of Experimental Marine Biology and Ecology 272: 131-146.
- Schroeder, W. W. 2002. Observations of *Lophelia pertusa* and the surficial geology at a deep-water site in the northeastern Gulf of Mexico. Hydrobiologia 471:29-33.
- Sheriff, R.E, 2002, Encyclopedic Dictionary of Applied Geophysics, fourth Addition: Society of Exploration Geophysicists, Talsa, Oklahoma, 429 p.
- Starck WAI, Colin PL (1978) *Gramma linki*: a new species of grammid fish from the tropical western Atlantic. *Bull Mar Sci* 28, 146-152.
- Sulak, K.J., Brooks, R.A., Luke, K.E., Norem, A.D., Randall, M., Quaid, A.J., Yeargin, G.E., Miller, J.M., Harden, W.M., Caruso, J.H., Ross, S.W., 2007. Demersal fishes associated with Lophelia pertusa coral and hard-substrate biotopes on the continental slope, northern Gulf of Mexico. Bulletin of Marine Science. 81:65-92
- Taylor MS, Hellberg ME (2003) Genetic evidence for local retention of pelagic larvae in a Caribbean reef fish. *Science* **299**, 107-109.

- Tissot, B. N., M. M. Yoklavich, M. S. Love, k. York, and M. Amend. 2006. Structure-forming invertebrates as components of benthic habitat on deep banks off southern California with special reference to deep sea corals. Fish. Bull. 104: 167–181
- Turley, C., Roberts, J.M. & Guinotte, J.J. (2007) Corals in deep water: Will the unseen hand of ocean acidification destroy cold water ecosystems? Coral Reefs 26: 445-448.
- Vardaro, M., I. R. MacDonald, L. C. Bender and N. L. Guinasso Jr. (2005). "Dynamic biological and physical processes observed at a gas hydrate outcropping on the continental slope of the Gulf of Mexico." Geo-Marine Letters 26(1): 6-15.
- Wakeley JW (2004) Metapopulation models for historical inference. *Molecular Ecology* 13, 865-875.
- Waller, RG & Tyler PA. (2005) The reproductive biology of two deep-sea, reef-building scleractinians from the NE Atlantic Ocean. Coral Reefs, 24(3), 514-522.
- Wang JL, Whitlock MC (2003) Estimating effective population size and migration rates from genetic samples over space and time. *Genetics* **163**, 429-446.
- Warren, Daniel. 2006a. Archaeological assessment Report, "Shallow Coulomb" Prospect, Block 701 and Vicinity, Mississippi Canyon Area. Prepared by C & C Technologies, Inc. for Shell Exploration and Production. Report submitted to the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Warren, Daniel. 2006b. Archaeological Assessment Report, Blocks 1007 and 1008, Ewing Banks Area and Blocks 37 and 38, Green Canyon Area. Prepared by C & C Technologies, Inc. for ATP Oil and Gas. Report submitted to the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA.
- Wiggins, Melanie. 1995. Torpedoes in the Gulf, Galveston and the U-boats, 1942-1943. Texas A&M Press: College Station, TX.
- Wilson JB (1979b) Patch development of the deep-water coral *Lophelia pertusa* (L.) on Rockall Bank. J. Mar. Biol. Ass. U.K. 59:165-177.
- Yoklavich, M. and Love, M. (2005). Christmas tree corals: a new species discovered off southern California. *Current, the Journal of Marine Education*, **21**, 27-30.